

# On Diophantine quintuples

ANDREJ DUJELLA (Zagreb, Croatia)

## 1 Introduction

The Greek mathematician Diophantus of Alexandria noted that the set  $\{\frac{1}{16}, \frac{33}{16}, \frac{17}{4}, \frac{105}{16}\}$  has the following property: the product of any two of its distinct elements increased by 1 is a square of a rational number (see [5]). Fermat first found a set of four positive integers with the above property, and it was  $\{1, 3, 8, 120\}$ .

Let  $n$  be an integer. A set of positive integers  $\{x_1, x_2, \dots, x_m\}$  is said to have *the property  $D(n)$*  if for all  $1 \leq i < j \leq m$  the following holds:  $x_i x_j + n = y_{ij}^2$ , where  $y_{ij}$  is an integer. Such a set is called *a Diophantine  $m$ -tuple*.

Davenport and Baker [4] showed that if  $d$  is a positive integer such that the set  $\{1, 3, 8, d\}$  has the property of Diophantus, then  $d$  has to be 120. This implies that the Diophantine quadruple  $\{1, 3, 8, 120\}$  cannot be extended to the Diophantine quintuple with the property  $D(1)$ . Analogous result was proved for the Diophantine quadruple  $\{2, 4, 12, 420\}$  with the property  $D(1)$  [17], for the Diophantine quadruple  $\{1, 5, 12, 96\}$  with the property  $D(4)$  [15] and for the Diophantine quadruples  $\{k-1, k+1, 4k, 16k^3-4k\}$  with the property  $D(1)$  for almost all positive integers  $k$  [9].

Euler proved that every Diophantine pair  $\{x_1, x_2\}$  with the property  $D(1)$  can be extended in infinitely many ways to the Diophantine quadruple with the same property (see [12]). In [6] it was proved that the same conclusion is valid for the pair with the property  $D(l^2)$  if the additional condition that  $x_1 x_2$  is not a perfect square is fulfilled.

Arkin, Hoggatt and Strauss [3] proved that every Diophantine triple with the property  $D(1)$  can be extended to the Diophantine quadruple. More precisely, if  $x_i x_j + 1 = y_{ij}^2$ , then we can set  $x_4 = x_1 + x_2 + x_3 + 2x_1 x_2 x_3 + 2y_{12} y_{13} y_{23}$ . For the Diophantine quadruple obtained in this way, they proved the existence of a positive rational number  $x_5$  with the property that  $x_i x_5 + 1$  is a square of a rational number for  $i = 1, 2, 3, 4$ .

Using this construction, in [2, 7, 8, 11] some formulas for Diophantine

quintuples in the terms of polynomials, Fibonacci, Lucas, Pell and Pell-Lucas numbers were obtained.

In the present paper we prove that for all positive rational numbers  $q, x_1, x_2, x_3, x_4$  such that  $x_i x_j + q^2 = y_{ij}^2$ ,  $y_{ij} \in \mathbf{Q}$ , for  $1 \leq i < j \leq 4$ , and  $x_1 x_2 x_3 x_4 \neq q^4$ , there exists a positive rational number  $x_5$  such that  $x_i x_5 + q^2$  is a square of a rational number for  $i = 1, 2, 3, 4$ . As a corollary we get the result that for *all* Diophantine quadruples  $\{x_1, x_2, x_3, x_4\}$  with the property  $D(1)$  there exists a rational number  $x_5$  such that  $x_i x_5 + 1$  is a square of a rational number for  $i = 1, 2, 3, 4$ .

## 2 Extension of Diophantine quadruples

**THEOREM 1** *Let  $q, x_1, x_2, x_3, x_4$  be rational numbers such that  $x_i x_j + q^2 = y_{ij}^2$ ,  $y_{ij} \in \mathbf{Q}$ , for all  $1 \leq i < j \leq 4$ . Assume that  $x_1 x_2 x_3 x_4 \neq q^4$ . Then the rational number  $x_5 = A/B$ , where*

$$\begin{aligned} A &= q^3[2y_{12}y_{13}y_{14}y_{23}y_{24}y_{34} + qx_1x_2x_3x_4(x_1 + x_2 + x_3 + x_4) \\ &\quad + 2q^3(x_1x_2x_3 + x_1x_2x_4 + x_1x_3x_4 + x_2x_3x_4) + q^5(x_1 + x_2 + x_3 + x_4)], \\ B &= (x_1x_2x_3x_4 - q^4)^2, \end{aligned}$$

has the property that  $x_i x_5 + q^2$  is a square of a rational number for  $i = 1, 2, 3, 4$ . To be more precise, for  $i \in \{1, 2, 3, 4\}$  it holds:

$$x_i x_5 + q^2 = \left( q \frac{x_i y_{jk} y_{jl} y_{kl} + q y_{ij} y_{ik} y_{il}}{x_1 x_2 x_3 x_4 - q^4} \right)^2,$$

where  $\{i, j, k, l\} = \{1, 2, 3, 4\}$ .

*Proof.* Let  $i \in \{1, 2, 3, 4\}$  and  $\{i, j, k, l\} = \{1, 2, 3, 4\}$ . Then we have:

$$\begin{aligned} &(x_1 x_2 x_3 x_4 - q^4)^2 (x_i x_5 + q^2) \\ &= 2q^3 x_i y_{12} y_{13} y_{14} y_{23} y_{24} y_{34} + q^4 x_1 x_2 x_3 x_4 x_i (x_1 + x_2 + x_3 + x_4) \\ &\quad + 2x_i q^6 (x_1 x_2 x_3 + x_1 x_2 x_4 + x_1 x_3 x_4 + x_2 x_3 x_4) \\ &\quad + x_i q^8 (x_1 + x_2 + x_3 + x_4) + q^2 x_1^2 x_2^2 x_3^2 x_4^2 - 2q^6 x_1 x_2 x_3 x_4 + q^{10} \\ &= q^2 [2q x_i y_{12} y_{13} y_{14} y_{23} y_{24} y_{34} + q^2 x_i^2 x_j x_k x_l (x_i + x_j + x_k + x_l) \\ &\quad + 2q^4 x_i^2 (x_j x_k + x_j x_l + x_k x_l) + 2q^4 x_i x_j x_k x_l + q^6 x_i^2 \\ &\quad + q^6 (x_i x_j + x_i x_k + x_i x_l) + x_i^2 x_j^2 x_k^2 x_l^2 - 2q^4 x_i x_j x_k x_l + q^8] \\ &= q^2 [2q x_i y_{12} y_{13} y_{14} y_{23} y_{24} y_{34} + x_i^2 (x_j x_k + q^2)(x_j x_l + q^2)(x_k x_l + q^2) \\ &\quad + q^2 (x_i x_j + q^2)(x_i x_k + q^2)(x_i x_l + q^2)] \\ &= q^2 (2q x_i y_{12} y_{13} y_{14} y_{23} y_{24} y_{34} + x_i^2 y_{jk}^2 y_{jl}^2 y_{kl}^2 + q^2 y_{ij}^2 y_{ik}^2 y_{il}^2) \\ &= [q(x_i y_{jk} y_{jl} y_{kl} + q y_{ij} y_{ik} y_{il})]^2, \end{aligned}$$

which proves the theorem.  $\blacksquare$

Since the signs of  $y_{ij}$  are arbitrary, we have two choices for  $x_5$ . Let  $x_5^+$  and  $x_5^-$  denote these two numbers, and let  $x_5^+$  be the number which corresponds to the case where all  $y_{ij}$  are nonnegative.

**COROLLARY 1** *Let  $\{x_1, x_2, x_3, x_4\} \subset \mathbf{N}$  be the set with the property  $D(1)$ . Then there exists a rational number  $x_5$ ,  $0 < x_5 < 1$ , such that  $x_i x_5 + 1$  is a square of a rational number for  $i = 1, 2, 3, 4$ .*

*Proof.* We claim that the number  $x_5^+$ , obtained by applying the construction from Theorem 1 to the set  $\{x_1, x_2, x_3, x_4\}$ , has the desired property. Indeed, it is sufficient to prove that  $x_5^+ < 1$ . Let us introduce the following notation:

$$\begin{aligned}\sigma_1 &= x_1 + x_2 + x_3 + x_4 \\ \sigma_2 &= x_1 x_2 + x_1 x_3 + x_1 x_4 + x_2 x_3 + x_2 x_4 + x_3 x_4 \\ \sigma_3 &= x_1 x_2 x_3 + x_1 x_2 x_4 + x_1 x_3 x_4 + x_2 x_3 x_4 \\ \sigma_4 &= x_1 x_2 x_3 x_4 \\ X &= \sigma_1 \sigma_4 + 2\sigma_3 + \sigma_1 \\ Y &= y_{12} y_{13} y_{14} y_{23} y_{24} y_{34}.\end{aligned}$$

The proof that  $x_5^+ = \frac{2Y + X}{(\sigma_4 - 1)^2} < 1$  is completed by showing that

$$2X < (\sigma_4 - 1)^2 \quad \text{and} \quad 4Y < (\sigma_4 - 1)^2. \quad (1)$$

Without loss of generality we can assume that  $x_1 < x_2 < x_3 < x_4$ . If  $x_1 = 1$ , then  $x_2 \neq 2$ . Therefore,  $x_2 \geq 3$ ,  $x_3 \geq 4$  and  $x_4 \geq 5$ . Hence  $\sigma_4 \geq 60$ . Furthermore, from

$$\frac{1}{x_1 x_2 x_3} + \frac{1}{x_1 x_2 x_4} + \frac{1}{x_1 x_3 x_4} + \frac{1}{x_2 x_3 x_4} \leq \frac{13}{60} < \frac{1}{4}$$

it follows that  $52 \leq 4\sigma_1 < \sigma_4$ . In the same manner we can see that  $59 \leq \sigma_2 < \sigma_4$  and  $107 \leq \sigma_3 < 2\sigma_4$  (see also [12]). Hence

$$(\sigma_4 - 1)^2 - 2X > \sigma_4^2 - 2\sigma_4 + 1 - \frac{\sigma_4^2}{2} - 8\sigma_4 - \frac{\sigma_4}{2} = \frac{1}{2}(\sigma_4^2 - 21\sigma_4 + 2) > 0$$

(since  $\sigma_4 \geq 60$ ). To get the second inequality from (1), we note that

$$Y^2 = (x_1 x_2 + 1)(x_1 x_3 + 1)(x_1 x_4 + 1)(x_2 x_3 + 1)(x_2 x_4 + 1)(x_3 x_4 + 1)$$

$$\begin{aligned}
&= \sigma_4^3 + \sigma_2\sigma_4^2 - \sigma_4^2 + \sigma_1\sigma_3\sigma_4 + \sigma_1^2\sigma_4 - 2\sigma_2\sigma_4 + \sigma_3^2 - \sigma_4 + \sigma_1\sigma_3 + \sigma_2 + 1 \\
&< \sigma_4^3 + \sigma_4^3 - \sigma_4^2 + \frac{\sigma_4^2}{2} + \frac{\sigma_4^2}{16} - 118\sigma_4 + 4\sigma_4^2 - \sigma_4 + \frac{\sigma_4^2}{2} + \sigma_4 + 1 \\
&= \frac{41}{16}\sigma_4^3 + \frac{7}{2}\sigma_4^2 - 118\sigma_4 + 1.
\end{aligned}$$

Therefore,

$$\begin{aligned}
&(\sigma_4 - 1)^4 - 16Y^2 \\
&> \sigma_4^4 - 4\sigma_4^3 + 6\sigma_4^2 - 4\sigma_4 + 1 - 41\sigma_4^3 - 56\sigma_4^2 + 1888\sigma_4 - 16 \\
&= \sigma_4^4 - 45\sigma_4^3 - 50\sigma_4^2 + 1884\sigma_4 - 15 > 0
\end{aligned}$$

(since  $\sigma_4 \geq 60$ ), which completes the proof.  $\blacksquare$

**COROLLARY 2** *Let  $q, x_1, x_2, x_3$  be rational numbers such that  $x_i x_j + q^2 = y_{ij}^2$ ,  $y_{ij} \in \mathbf{Q}$  for all  $1 \leq i < j \leq 3$ . Let*

$$\begin{aligned}
x_4 &= [2y_{12}y_{13}y_{23} + 2x_1x_2x_3 + q^2(x_1 + x_2 + x_3)]/q^2, \\
x_5 &= \frac{4y_{12}y_{13}y_{23}(x_1y_{23} + y_{12}y_{13})(x_2y_{13} + y_{12}y_{23})(x_3y_{12} + y_{13}y_{23})}{(x_1x_2x_3x_4 - q^4)^2}.
\end{aligned}$$

*Then the set  $\{x_1, x_2, x_3, x_4, x_5\}$  has the property that the product of its any two distinct elements increased by  $q^2$  is equal to the square of a rational number. In the notation of Theorem 1, we have*

$$x_5 = \frac{4q^3 y_{12} y_{13} y_{14} y_{23} y_{24} y_{34}}{(x_1 x_2 x_3 x_4 - q^4)^2}.$$

*Proof.* Let  $z_1 = x_1, z_2 = x_2, z_3 = x_3, z_4 = 0$ . Then the rational numbers  $z_1, z_2, z_3, z_4$  satisfy the conditions of Theorem 1, and its application gives us the number

$$z_5 = [2y_{12}y_{13}y_{23} + 2x_1x_2x_3 + q^2(x_1 + x_2 + x_3)]/q^2.$$

Set  $x_4 = z_5$ . We can now apply Theorem 1 on the numbers  $x_1, x_2, x_3, x_4$ . Let  $x_5$  be the number which is obtained by this construction. Observe that, by Theorem 1, for all  $i \in \{1, 2, 3\}$

$$qy_{i4} = x_i y_{jk} + y_{ij} y_{ik}, \quad (2)$$

where  $\{i, j, k\} = \{1, 2, 3\}$ . Let us introduce the following notation:

$$\begin{aligned}\Sigma_1 &= x_1 + x_2 + x_3 \\ \Sigma_2 &= x_1x_2 + x_1x_3 + x_2x_3 \\ \Sigma_3 &= x_1x_2x_3 \\ V &= y_{12}y_{13}y_{23} \\ W &= y_{14}y_{24}y_{34}.\end{aligned}$$

We have

$$V^2 = (x_1x_2 + q^2)(x_1x_3 + q^2)(x_2x_3 + q^2) = \Sigma_3^2 + q^2\Sigma_1\Sigma_3 + q^4\Sigma_2 + q^6.$$

From (2) it follows that

$$\begin{aligned}q^3W &= (x_1y_{23} + y_{12}y_{13})(x_2y_{13} + y_{12}y_{23})(x_3y_{12} + y_{13}y_{23}) \\ &= 4\Sigma_3^2 + 3q^2\Sigma_1\Sigma_3 + 2q^4\Sigma_2 + q^6 + V(4\Sigma_3 + q^2\Sigma_1).\end{aligned}$$

Now it is easy to check that, in notation of Corollary 1,

$$q^4\sigma_1\sigma_4 + 2q^6\sigma_3 + q^8\sigma_1 = 2q^3VW. \quad (3)$$

Consequently,

$$\begin{aligned}x_5 &= \frac{4q^3VW}{(x_1x_2x_3x_4 - q^4)^2} \\ &= \frac{4q^3y_{12}y_{13}y_{14}y_{23}y_{24}y_{34}}{(x_1x_2x_3x_4 - q^4)^2} \\ &= \frac{4y_{12}y_{13}y_{23}(x_1y_{23} + y_{12}y_{13})(x_2y_{13} + y_{12}y_{23})(x_3y_{12} + y_{13}y_{23})}{(x_1x_2x_3x_4 - q^4)^2}.\end{aligned}$$

■

Let us now consider the question when one or both (since  $x_5^+$  and  $x_5^-$  can be equal) of the numbers  $x_5^+$  and  $x_5^-$  will be equal to zero. For the obvious reason, such extension of a Diophantine quadruple we will call trivial. We will see that the answer to this question is closely connected to the construction of Corollary 2. From now on, we assume that  $q \neq 0$ .

**PROPOSITION 1** *In the notation of Theorem 1, we have  $x_5^+ = x_5^- = 0$  if and only if there exist  $1 \leq i < j \leq 4$  such that  $x_ix_j = -q^2$  and  $x_i + x_j = x_k + x_l$ , where  $\{i, j, k, l\} = \{1, 2, 3, 4\}$ .*

*Proof.* From  $x_5^+ = x_5^-$  we conclude that there exist  $1 \leq i < j \leq 4$  such that  $y_{ij} = 0$ , i.e.  $x_i x_j = -q^2$ . Substituting this into expression for  $x_5$  we obtain

$$x_5 = \frac{q^2(x_i + x_j - x_k - x_l)}{x_k x_l + q^2}. \quad (4)$$

Consequently, the condition  $x_5 = 0$  implies that  $x_i + x_j = x_k + x_l$ .

Conversely, suppose that  $x_1, x_2, x_3, x_4$  satisfy the condition of the proposition. Then  $y_{ij} = 0$ , and (4) implies that  $x_5^+ = x_5^- = 0$ .  $\blacksquare$

**PROPOSITION 2** *In the notation of Theorem 1, we have  $0 \in \{x_5^+, x_5^-\}$  if and only if there exists  $i \in \{1, 2, 3, 4\}$  such that*

$$x_i = [2y_{jk}y_{jl}y_{kl} + 2x_j x_k x_l + q^2(x_j + x_k + x_l)]/q^2,$$

where  $\{i, j, k, l\} = \{1, 2, 3, 4\}$ .

*Proof.* We can assume that  $y_{ij} \neq 0$  since otherwise the assertion of the proposition follows from Proposition 1. If  $x_5 = 0$ , then  $x_i x_5 + q^2 = q^2$  for  $i = 1, 2, 3, 4$ . Hence, if  $0 \in \{x_5^+, x_5^-\}$ , then Theorem 1 implies that for appropriate choice of the sign of  $y_{ij}$  we have

$$x_i y_{jk} y_{jl} y_{kl} + q y_{ij} y_{ik} y_{il} = \pm(x_1 x_2 x_3 x_4 - q^4),$$

where  $\{i, j, k, l\} = \{1, 2, 3, 4\}$ . Hence, there is no loss of generality is assuming that

$$x_1 y_{23} y_{24} y_{34} + q y_{12} y_{13} y_{14} = x_2 y_{13} y_{14} y_{34} + q y_{12} y_{23} y_{24}.$$

This gives  $(x_1 y_{34} - q y_{12}) y_{23} y_{24} = (x_2 y_{34} - q y_{12}) y_{13} y_{14}$ . Set  $x_1 y_{34} - q y_{12} = \alpha y_{13} y_{14}$ . Then  $x_2 y_{34} - q y_{12} = \alpha y_{23} y_{24}$ , and so

$$\begin{aligned} & \alpha(x_1 y_{23} y_{24} y_{34} + q y_{12} y_{13} y_{14}) \\ &= x_1 y_{34} (x_2 y_{34} - q y_{12}) + q y_{12} (x_1 y_{34} - q y_{12}) \\ &= x_1 x_2 y_{34}^2 - q^2 y_{12}^2 = x_1 x_2 x_3 x_4 - q^4. \end{aligned}$$

We thus get  $\alpha = \pm 1$  and  $x_1 y_{34} - q y_{12} = \pm y_{13} y_{14}$ . Squaring this relation we obtain

$$x_1^2 x_3 x_4 + q^2 x_1^2 + q^2 x_1 x_2 + q^4 - 2q x_1 y_{12} y_{34} = x_1^2 x_3 x_4 + q^2 x_1 x_3 + q^2 x_1 x_4 + q^4,$$

and (if  $x_1 \neq 0$ )  $2y_{12}y_{34} = q(x_1 + x_2 - x_3 - x_4)$ . Squaring again we obtain the quadratic equation in  $x_4$ :

$$q^2x_4^2 - 2x_4[q^2(x_1 + x_2 + x_3) + 2x_1x_2x_3] + q^2(x_1^2 + x_2^2 + x_3^2 - 2x_1x_2 - 2x_1x_3 - 2x_2x_3 - 4q^2) = 0,$$

with the solutions

$$x_4 = [q^2(x_1 + x_2 + x_3) + 2x_1x_2x_3 \pm 2y_{12}y_{13}y_{23}]/q^2. \quad (5)$$

We have been working under assumption that  $x_1 \neq 0$ . Now suppose that  $x_1 = 0$ . In the same manner, using Corollary 2, it can be proved that  $x_1 = 0$  implies

$$x_4 = x_2 + x_3 \pm y_{23},$$

which is exactly the relation (5) for  $x_1 = 0$ .

This proves one implication of the proposition. The opposite implication is direct consequence of the relation (3). ■

### 3 Examples

EXAMPLE 1 Let us first show that the condition  $x_1x_2x_3x_4 \neq q^4$  from Theorem 1 is not superfluous. Indeed, the set  $\{25600, 50625, 82944, 518400\}$  has the property  $D(86400^2)$  and

$$25600 \cdot 50625 \cdot 82944 \cdot 518400 = 86400^4.$$

As an illustration of the situation from Proposition 1 let us adduce the set  $\{-25, 25, -24, 24\}$  with the property  $D(625)$  and the set  $\{-1, 64, 48, 15\}$  with the property  $D(64)$ . In both cases the construction from Theorem 1 gives  $x_5^+ = x_5^- = 0$ .

From [10, (13)], for  $a = 2$  and  $k = 3$ , we obtain the Diophantine quadruple  $\{2, 20, 44, 72\}$  with the property  $D(81)$ . It is easy to check that this quadruple does not satisfy the conditions of Proposition 2. Therefore the numbers  $x_5^+$  and  $x_5^-$  are different from 0. Indeed,  $x_5^+ = \frac{4860}{169}$  and  $x_5^- = -\frac{1156680}{1054729}$ . Using  $x_5^+$ , we obtain the Diophantine quintuple  $\{338, 3380, 4860, 7436, 12168\}$  with the property  $D(39^4)$ .

If we apply the construction from Theorem 1 to the original set of Diophantus  $\{\frac{1}{16}, \frac{33}{16}, \frac{17}{4}, \frac{105}{16}\}$ , we obtain  $x_5^+ = \frac{549120}{101^2}$  and  $x_5^- = \frac{-26880}{421^2}$ .

The definition of a Diophantine  $m$ -tuple can be extended to the subsets of  $\mathbf{Q}$ . Let  $q$  be a rational number. We call a set  $A = \{x_1, x_2, \dots, x_m\} \subset$

$\mathbf{Q} \setminus \{0\}$  a (rational) Diophantine  $m$ -tuple with the property  $D(q)$  if the product of any two distinct elements of  $A$  increased by  $q$  is equal to the square of a rational number. The construction of the rational Diophantine quintuple with the property  $D(1)$  which extends the given Diophantine triple was described in [3]. That construction is equivalent to the construction from Corollary 2. But Theorem 1 makes possible the extension of the Diophantine quadruples which are not of the form  $\{x_1, x_2, x_3, x_4\}$  from Corollary 2. One such quadruple is the set  $\{2, 20, 44, 72\}$  from Example 1. Let us now examine two ways for generation of such Diophantine quadruples.

EXAMPLE 2 Let  $\{x_1, x_2, x_3, x_4\} \subset \mathbf{Q}$  be an arbitrary set with the property  $D(q^2)$  and let  $x_5 \in \mathbf{Q}$  be the number which is obtained by applying Theorem 1 to this set. Then the set  $\{x_2, x_3, x_4, x_5\}$  also has the property  $D(q^2)$ , and we can apply Theorem 1 again. In this way we obtain  $x_6 \in \mathbf{Q}$  such that the set  $\{x_2, x_3, x_4, x_5, x_6\}$  has the property  $D(q^2)$ .

For example, if  $x_1 = k - 1$ ,  $x_2 = k + 1$ ,  $x_3 = 4k$  and  $x_4 = 16k^3 - 4k$ , then the set  $\{x_1, x_2, x_3, x_4\}$  has the property  $D(1)$  ([6, p. 22]) and we obtain

$$x_5 = \frac{4k(2k-1)(2k+1)(4k^2-2k-1)(4k^2+2k-1)(8k^2-1)}{(64k^6-80k^4+16k^2-1)^2},$$

and  $x_6 = P(k)/Q(k)$ , where

$$\begin{aligned} P(k) &= (8k^3 - 4k^2 + 1)(8k^3 + 4k^2 - 4k - 1)(8k^3 - 12k^2 + 1) \\ &\quad \times (8k^4 + 4k^3 - 8k^2 - k + 1)(32k^4 - 8k^3 + 28k^2 + 3) \\ &\quad \times (32k^4 + 8k^3 - 12k^2 + 1)(32k^4 + 24k^3 - 12k^2 - 4k + 1) \\ &\quad \times (32k^4 + 40k^3 + 4k^2 - 4k + 1), \end{aligned}$$

$$\begin{aligned} Q(k) &= (131072k^{14} + 131072k^{13} - 184320k^{12} - 180224k^{11} + 96256k^{10} \\ &\quad + 86016k^9 - 26880k^8 - 18432k^7 + 4480k^6 + 1792k^5 - 480k^4 \\ &\quad - 64k^3 + 32k^2 - 1)^2. \end{aligned}$$

It turns out that this factorization of the numerator of  $x_6$  is not accidental. Namely, it can be checked that, in notation of Theorem 1,  $x_6 = P/Q$ , where

$$\begin{aligned} P &= q^3(y_{12}y_{13}y_{14} + qy_{12}y_{13} + qy_{12}y_{23} + qy_{13}y_{23}) \\ &\quad \times (y_{12}y_{13}y_{14} + qy_{12}y_{13} - qy_{12}y_{23} - qy_{13}y_{23}) \\ &\quad \times (y_{12}y_{13}y_{14} - qy_{12}y_{13} + qy_{12}y_{23} - qy_{13}y_{23}) \\ &\quad \times (y_{12}y_{13}y_{14} - qy_{12}y_{13} - qy_{12}y_{23} + qy_{13}y_{23})(y_{23}y_{24} + y_{23}y_{34} + y_{24}y_{34}) \\ &\quad \times (y_{23}y_{24} + y_{23}y_{34} - y_{24}y_{34})(y_{23}y_{24} - y_{23}y_{34} + y_{24}y_{34}) \\ &\quad \times (-y_{23}y_{24} + y_{23}y_{34} + y_{24}y_{34}), \end{aligned}$$



$$Q = x_1^4(4x_2x_3x_4y_{12}y_{13}y_{14}y_{23}y_{24}y_{34} - qx_1^2x_2^2x_3^2x_4^2 + 2q^5x_1x_2x_3x_4 - q^9)^2.$$

PROPOSITION 3 *Let  $x_1$ ,  $x_2$ , and  $x_3$  be rational numbers such that the denominator of*

$$x_4 = \frac{8(x_3 - x_1 - x_2)(x_1 + x_3 - x_2)(x_2 + x_3 - x_1)}{(x_1^2 + x_2^2 + x_3^2 - 2x_1x_2 - 2x_1x_3 - 2x_2x_3)^2}$$

*is different from 0. Then  $x_1x_4 + 1$ ,  $x_2x_4 + 1$  and  $x_3x_4 + 1$  are squares of rational numbers.*

*Proof.* It follows immediately that

$$x_1x_4 + 1 = \left( \frac{x_2^2 - 2x_2x_3 + x_3^2 - 3x_1^2 + 2x_1x_2 + 2x_1x_3}{x_1^2 + x_2^2 + x_3^2 - 2x_1x_2 - 2x_1x_3 - 2x_2x_3} \right)^2,$$

and analogous relations hold for  $x_2x_4 + 1$  and  $x_3x_4 + 1$ . ■

EXAMPLE 3 Let us observe that the set  $\{x_1, x_2, x_3\}$  in Proposition 3 does not need to have the property  $D(1)$ . Let us take for example  $x_1 = F_{2n+1}$ ,  $x_2 = F_{2n+3}$  and  $x_3 = F_{2n+5}$ . Then the set  $\{x_1, x_2, x_3\}$  has the property  $D(-1)$  for every positive integer  $n$  (see [13, 14]). Proposition 3 implies that there exists a rational number  $x_4$  with the property that  $x_ix_4 + 1$ ,  $i = 1, 2, 3$ , are squares of rational numbers. We will show that in this case the number  $x_4$  is an integer. Indeed,

$$\begin{aligned} x_1^2 + x_2^2 + x_3^2 - 2x_1x_2 - 2x_1x_3 - 2x_2x_3 &= (x_1 - x_2 + x_3)^2 - 4x_1x_3 \\ &= [F_{2n+1} - F_{2n+3} + (3F_{2n+3} - F_{2n+1})]^2 - 4F_{2n+1}F_{2n+5} \\ &= 4(F_{2n+3}^2 - F_{2n+1}F_{2n+5}) = -4. \end{aligned}$$

Hence,

$$x_4 = \frac{8}{16} \cdot 2F_{2n+2} \cdot 2F_{2n+3} \cdot 2F_{2n+4} = 4F_{2n+2}F_{2n+3}F_{2n+4}.$$

EXAMPLE 4 If  $x_1x_2 + 1 = y_{12}^2$  and  $x_3 = x_1 + x_2 + 2y_{12}$ , then the set  $\{x_1, x_2, x_3\}$  has the property  $D(1)$ . If we apply the construction from Proposition 3 to this set we obtain

$$x_4 = 4y_{12}(x_1 + y_{12})(x_2 + y_{12}).$$

If we apply the construction from Corollary 2 to the set  $\{x_1, x_2, x_3\}$  we obtain exactly the same result.

EXAMPLE 5 Let  $x_1 = 1$ ,  $x_2 = 3$  and  $x_3 = 120$ . Then proposition 3 gives  $x_4 = \frac{834968}{3361^2}$ . The set  $\{x_1, x_2, x_3, x_4\}$  has the property  $D(1)$  and we can apply the construction from Theorem 1. We obtain:

$$x_5^+ = \frac{3985166705520 \cdot 481^2}{601439^2 \cdot 481^2}, \quad x_5^- = \frac{426360 \cdot 601439^2}{481^2 \cdot 601439^2}.$$

It turns out that this cancelation is not accidental. Namely, let  $\{x_1, x_2, x_3\}$  be the arbitrary set with the property  $D(1)$ , let  $x_4$  be the number which is obtained by applying Proposition 3 to this set, and let  $x_5^+$  and  $x_5^-$  be the numbers which are obtained by applying Theorem 1 to the set  $\{x_1, x_2, x_3, x_4\}$ . Then

$$\sqrt{x_1 x_5^+ + 1} \cdot \sqrt{x_1 x_5^- + 1} = \left| \frac{(a+b)(a-b)cd}{c^2 d^2} \right|,$$

where

$$\begin{aligned} a &= x_1 y_{23} [x_1^2 (4x_2 x_3 + 1) - 2x_1 (x_2 + x_3) (2x_2 x_3 - 1) - (3x_2^2 + 2x_2 x_3 + 3x_3^2)], \\ b &= y_{12} y_{13} [x_1^2 (-4x_2 x_3 - 3) + 2x_1 (x_2 + x_3) (2x_2 x_3 + 1) + (x_2 - x_3)^2], \\ c &= (x_1 + x_2 + x_3)^2 - 4(x_1 x_2 x_3 - y_{12} y_{13} y_{23})^2 + 4, \\ d &= 4(x_1 x_2 x_3 + y_{12} y_{13} y_{23})^2 - (x_1 + x_2 + x_3)^2 - 4. \end{aligned}$$

For  $x_1 = 1$ ,  $x_2 = 3$  and  $x_3 = 120$ , we get  $c = 4 \cdot 481$  and  $d = 4 \cdot 601439$ .

## 4 Some open problems

One question still unanswered is whether there exists a (positive integer) Diophantine quintuple with the property  $D(1)$ . Corollary 1 shows that if such a quintuple exists it cannot be obtained by the construction from Theorem 1. Let us mention that the analogous result for the sets with the property  $D(l^2)$ , where  $l > 1$ , does not hold. For example, if we apply the construction from Theorem 1 to the quadruples  $\{4, 21, 69, 125\}$  and  $\{7, 12, 63, 128\}$  with the property  $D(400)$ , we obtain  $x_5^+ = 384$ ,  $x_5^- = -\frac{4032000}{1129^2}$  and  $x_5^+ = 375$ ,  $x_5^- = -\frac{11856000}{2021^2}$ , respectively. Hence, the sets  $\{4, 21, 69, 125, 384\}$  and  $\{7, 12, 69, 125, 375\}$  are Diophantine quintuples with the property  $D(400)$ .

One may ask which is the least positive integer  $n_1$ , and which is the greatest negative integer  $n_2$ , for which there exists a Diophantine quintuple with the property  $D(n_i)$ ,  $i = 1, 2$ . Certainly  $n_1 \leq 256$  and  $n_2 \geq -255$ , since the sets  $\{1, 33, 105, 320, 18240\}$  and  $\{5, 21, 64, 285, 6720\}$  have the property  $D(256)$ , and the set  $\{8, 32, 77, 203, 528\}$  has the property  $D(-255)$ .

In present paper we have considered the quintuples with the property  $D(q)$ , where  $q$  was a square of a rational number. However, the last set with the property  $D(-255)$  indicates that there exist quintuples with the property  $D(q)$ , where  $q$  is not a perfect square (see also [9, 16]). Thus we came to the following open problem: For which rational numbers  $q$  there exists a rational Diophantine quintuple with the property  $D(q)$ ? It follows easily from [6, Theorem 5] that for every rational number  $q$  there exists a rational Diophantine quadruple with the property  $D(q)$ .

At present it is not known whether there exists a rational number  $q \neq 0$  such that there exists a rational Diophantine sextuple with the property  $D(q)$ . In [1], some rational "sextuples" with the property  $D(1)$  were obtained, but all of them have two equal elements. Thus, they are actually quintuples with the additional property that  $x_1^2 + 1$  is a perfect square. There exists also a rational Diophantine quintuple  $\{x_1, \dots, x_5\}$  with the property  $D(1)$  such that  $x_1^2 + 1$ ,  $x_2^2 + 1$  and  $x_3^2 + 1$  are perfect squares. However, the question of the existence of Diophantine sextuples is still open.

## References

- [1] J. Arkin, D. C. Arney, F. R. Giordano, R. A. Kolb and G. E. Bergum, *An extension of an old classical Diophantine problem*, in: Application of Fibonacci Numbers, Vol. 5, G. E. Bergum, A. N. Philippou and A. F. Horadam (eds.), Kluwer, Dordrecht, 1993, 45-48.
- [2] J. Arkin and G. E. Bergum, *More on the problem of Diophantus*, in: Application of Fibonacci Numbers, Vol. 2, A. N. Philippou, A. F. Horadam and G. E. Bergum (eds.), Kluwer, Dordrecht, 1988, 177-181.
- [3] J. Arkin, V. E. Hoggatt and E. G. Strauss, *On Euler's solution of a problem of Diophantus*, Fibonacci Quart. 17 (1979), 333-339.
- [4] H. Davenport and A. Baker, *The equations  $3x^2 - 2 = y^2$  and  $8x^2 - 7 = z^2$* , Quart. J. Math. Oxford Ser. (2) 20 (1969), 129-137.
- [5] Diophantus of Alexandria, *Arithmetics and the Book of Polygonal Numbers*, Nauka, Moscow, 1974 (in Russian).
- [6] A. Dujella, *Generalization of a problem of Diophantus*, Acta Arith. 65 (1993), 15-27.
- [7] A. Dujella, *Diophantine quadruples for squares of Fibonacci and Lucas numbers*, Portugal. Math. 52 (1995), 305-318.

- [8] A. Dujella, *Generalized Fibonacci numbers and the problem of Diophantus*, Fibonacci Quart. 34 (1996), 164-175.
- [9] A. Dujella, *Generalization of the Problem of Diophantus and Davenport*. Dissertation, University of Zagreb, 1996 (in Croatian).
- [10] A. Dujella, *Some polynomial formulas for Diophantine quadruples*, Grazer Math. Ber. (to appear).
- [11] A. Dujella, *A problem of Diophantus and Pell numbers*, preprint.
- [12] P. Heichelheim, *The study of positive integers  $(a, b)$  such that  $ab + 1$  is a square*, Fibonacci Quart. 17 (1979), 269-274.
- [13] V. E. Hoggatt and G. E. Bergum, *A problem of Fermat and the Fibonacci sequence*, Fibonacci Quart. 15 (1977), 323-330.
- [14] C. Long, G. E. Bergum, *On a problem of Diophantus*, in: Application of Fibonacci Numbers, Vol. 2, A. N. Philippou, A. F. Horadam and G. E. Bergum (eds.), Kluwer, Dordrecht, 1988, 183-191.
- [15] S. P. Mohanty and M. S. Ramasamy, *The characteristic number of two simultaneous Pell's equations and its application*, Simon Stevin 59 (1985), 203-214.
- [16] V. K. Mootha, *On the set of numbers  $\{14, 22, 30, 42, 90\}$* , Acta Arith. 71 (1995), 259-263.
- [17] M. Velupillai, *The equations  $z^2 - 3y^2 = -2$  and  $z^2 - 6x^2 = -5$* , in: A Collection of Manuscripts Related to the Fibonacci sequence, V. E. Hoggatt and M. Bicknell-Johnson (eds.), The Fibonacci Association, Santa Clara, 1980, 71-75.

DEPARTMENT OF MATHEMATICS  
UNIVERSITY OF ZAGREB  
BIJENIČKA CESTA 30  
10000 ZAGREB  
CROATIA  
E-mail: DUJE@MATH.HR